

On the Flatness and Injectivity of Dual Modules (II) *

HUANG Zhao-yong¹, TANG Jin-yu²

(1. Dept. of Math., Nanjing University, Jiangsu 210093, China;

2. Dept. of Math. & Info. Sci., Guangxi University, Nanning 530004, China)

Abstract: For a commutative ring R and an injective cogenerator E in the category of R -modules, we characterize QF rings, IF rings and semihereditary rings by using the properties of the dual modules with respect to E .

Key words: QF rings; flatness; injectivity; dual modules.

Classification: AMS(1991) 13D05,16E10/CLC O153.3

Document code: A **Article ID:** 1000-341X(2001)03-0377-07

1. Introduction

Throughout this paper, R will denote an associate, commutative ring with identity and all modules are unital. E always denotes a certain injective cogenerator in the category of R -modules.

Let M be an R -module. In [4] we introduce the notion of the dual module $\text{Hom}_R(M, E)$ with respect to E , and denote it by M^e . It is shown that the flatness of M^e is equivalent to the FP-injectivity or the injectivity of M if and only if R is a coherent ring or a noether ring respectively. The FP-injectivity, the injectivity of M and the projectivity of M^e are pairwise equivalent if and only if R is an artin ring (see [4]).

Recall that R is called a QF ring if R is an artin ring and for each ideal I of R , $I = 0 :_R (0 :_R I)$ (see [5]). Such rings have been extensively studied, many properties equivalent to this definition have been obtained. For example, the following statements are equivalent:

- (1) R is a QF ring;
- (2) R is a noether ring and for each ideal I of R , $I = 0 :_R (0 :_R I)$;
- (3) R is an artin ring (or a noether ring) and R is a cogenerator in the category of R -modules;

*Received date: 1998-10-26

Foundation item: Supported by National Natural Science Foundation of China (10001017) and Scientific Research Foundation for Returned Overseas Chinese Scholars (State Education Commission)

Biography: HUANG Zhao-yong (1968-), male, born in Daye county, Hubei province, Ph.D, associate professor.

E-mail: huangzy@netra.nju.edu.cn

- (4) R is an artin ring (or a noether ring) and R is selfinjective;
- (5) Any projective R -module is injective;
- (6) Any injective R -module is projective.

Also recall that R is called an IF ring if every injective R -module is flat (see [7]). It is shown that R is an IF ring if and only if R is a coherent ring and R is self FP-injective. It is clear that the notion of IF rings is a generalization of that of QF rings and Von Neumann regular rings.

In this paper we will introduce the notions of E -artin rings, E -coherent rings and f -cogenerators, and characterize QF rings, IF rings and semihereditary rings by using the properties of dual modules.

2. Main results

Proposition 1 *The following statements are equivalent.*

- (1) R is a QF ring;
- (2) R is a noether ring and R^e is flat;
- (3) R is an artin ring and R^e is flat;
- (4) M^e is a projective module for any flat module M ;
- (4)' M^e is a projective module for any projective module M ;
- (4)'' M^e is a projective module for any free module M ;
- (5) M^e is a submodule of a projective module for any flat module M ;
- (5)' M^e is a submodule of a projective module for any projective module M ;
- (5)'' M^e is a submodule of a projective module for any free module M .

Proof (2) \Leftrightarrow (1) Suppose that R is a noether ring and R^e is flat. Then R is an FP-injective R -module by [4, Corollary 2]. So R is selfinjective and hence R is a QF ring. The converse implication is trivial.

(1) \Rightarrow (3) Suppose R is a QF ring. Then R is an artin ring and any injective R -module is projective. Because $R^e \cong E$ is injective, R^e is projective.

(3) \Rightarrow (4) Suppose M is a flat module. Then there is a free module $R^{(I)}$ where I is a set such that $R^{(I)} \rightarrow M \rightarrow 0$ is exact. Since M is flat, this exact sequence is pure. So $0 \rightarrow M^e \rightarrow [R^{(I)}]^e \cong (R^e)^I$ is exact and splits by [4, Lemma 1]. It follows that M^e is a direct summand of $(R^e)^I$. Since R is an artin ring and R^e is flat, R^e is projective. So $(R^e)^I$ is also projective, it follows that M^e is projective.

(4) \Rightarrow (4)' \Rightarrow (4)'' \Rightarrow (5)'' and (4) \Rightarrow (5) \Rightarrow (5)' \Rightarrow (5)'' are trivial.

(5)'' \Rightarrow (1) Suppose M is an injective R -module. There is a free module F such that $F \rightarrow M^e \rightarrow 0$ is exact, so $0 \rightarrow M^{ee} \rightarrow F^e$ is exact. By (5)'' F^e is a submodule of a projective module P . It is known [4, Corollary 1] that M is a submodule of M^{ee} , we get that M is isomorphic to a submodule of P . So M is projective and R is a QF ring. \square

We know from Proposition 1 that R is a QF ring if and only if P^e is (a submodule of) a projective module for any projective module P . It is natural to ask what properties R possesses if Q^e is (a submodule of) a projective module for any injective module Q . [4, Theorem 3] says that R is an artin ring if and only if Q^e is projective for any injective module Q .

Definition 1 R is called an E -artin ring if Q^e is a submodule of a projective module for any injective module Q .

Remark 1 An artin ring is an E -artin ring by [4, Theorem 3]. If R is a hereditary ring, then R is artinian if and only if R is E -artinian.

Recall that an R -module A is called FP-injective if $\text{Ext}_R^1(B, A) = 0$ for any finitely presented R -module B . A ring R is called self FP-injective if R is FP-injective as an R -module (see [7]).

Lemma 1 Let R be an E -artin ring. Then the direct product of a family of projective modules can be embedded in a projective module.

Proof Suppose $\{P_i\}_{i \in I}$ is a family of projective modules where I is a set. Then each P_i^e is injective. By [3, Corollary 2.1.12] $\bigoplus_{i \in I} P_i^e$ is FP-injective, which implies that the exact sequence $0 \rightarrow \bigoplus_{i \in I} P_i^e \rightarrow \prod_{i \in I} P_i^e$ is pure, and so $(\prod_{i \in I} P_i^e)^e \rightarrow (\bigoplus_{i \in I} P_i^e)^e \cong \prod_{i \in I} P_i^{ee} \rightarrow 0$ splits. It follows that $\prod_{i \in I} P_i^{ee}$ is a direct summand of $(\prod_{i \in I} P_i^e)^e$. Because $\prod_{i \in I} P_i^e$ is injective and R is an E -artin ring, $(\prod_{i \in I} P_i^e)^e$ is a submodule of a projective module P . Since $\prod_{i \in I} P_i$ is a submodule of $\prod_{i \in I} P_i^{ee}$, $\prod_{i \in I} P_i$ can be embedded in P . \square

Lemma 2 The following statements are equivalent.

- (1) R is an E -artin ring;
- (2) M^e is a submodule of a projective module for any FP-injective module M ;
- (3) $\text{Hom}_R(B, C)$ is a submodule of a projective module for any injective module (or FP-injective module) B and any injective module C ;
- (4) P^{ee} is a submodule of a projective module for any flat module P ;
- (4)' P^{ee} is a submodule of a projective module for any projective module P ;
- (4)'' P^{ee} is a submodule of a projective module for any free module P .

Proof (1) \Rightarrow (2) Suppose M is an FP-injective module. Then the exact sequence $0 \rightarrow M \rightarrow E(M)$ is pure where $E(M)$ is the injective envelope of M . It follows from [4, Lemma 1] that M^e is a direct summand of $[E(M)]^e$. We know from (1) and Definition 1 that M^e is a submodule of a projective module.

(2) \Rightarrow (3) Suppose B is an FP-injective module and C is an injective module. Since E is an injective cogenerator in the category of R -modules, C is a direct summand of E^I for some set I . So $\text{Hom}_R(B, C)$ is a direct summand of $\text{Hom}_R(B, E^I) \cong (B^e)^I$. Since B^e is a submodule of a projective module P_1 by (2), $(B^e)^I$ is a submodule of P_1^I . It is known from Lemma 1 that P_1^I can be embedded in some projective module, and we are done.

(3) \Rightarrow (4) If P is a flat module, then P^e is injective. By (3) P^{ee} is a submodule of a projective module.

(4) \Rightarrow (4)' \Rightarrow (4)'' are trivial.

(4)'' \Rightarrow (1) For any injective module Q , Q is a direct summand of E^I for some set I , so Q^e is a direct summand of $(E^I)^e \cong [R^{(I)}]^{ee}$. By (4)'' Q^e is a submodule of a projective module, it follows that R is an E -artin ring. \square

Remark 2 Suppose both E and E' are injective cogenerators in the category of R -modules. By Lemma 2, R is an E -artin ring if and only if R is an E' -artin ring.

Definition 2 An R -module C is called an f -cogenerator in the category of R -modules if C cogenerates every finitely presented R -module.

Lemma 3 Let C be an f -cogenerator in the category of R -modules. Then

$$0 :_R (0 :_C I) = I$$

for any finitely generated ideal I of R .

Proof It is clear that $I \subseteq 0 :_R (0 :_C I)$. We only need to prove that

$$0 :_R (0 :_C I) \subseteq I.$$

Let $s \in R - I$. Since C is an f -cogenerator and R/I is finitely presented, C cogenerates R/I . Then we have a nonzero homomorphism $h : R/I \rightarrow C$ such that $h(s + I) \neq 0$. Suppose that $g : R \rightarrow R/I$ is the natural epimorphism. Then

$$0 = hg(I) = hg(1)I.$$

So $hg(1) \in 0 :_C I$. But

$$hg(1)s = hg(s) = h(s + I) \neq 0.$$

It follows that $s \notin 0 :_R (0 :_C I)$. So

$$0 :_R (0 :_C I) \subseteq I. \quad \square$$

For any R -modules M and N , recall from [1, p. 109] that

$$\text{Rej}_M(N) = \bigcap \{ \text{Ker } f \mid f \in \text{Hom}_R(M, N) \}.$$

Lemma 4 Let R be a coherent ring. Then R is an f -cogenerator in the category of R -modules if and only if R is self FP-injective.

Proof (\Rightarrow) Suppose that both I_1 and I_2 are finitely generated ideals of R . Since R is a coherent ring, it follows from [3, Theorem 2.3.2] that $I_1 \cap I_2$, $0 :_R I_1$ and $0 :_R I_2$ are finitely generated ideals of R . Then from Lemma 3 we get that

$$\begin{aligned} 0 :_R (0 :_R (I_1 \cap I_2)) &= I_1 \cap I_2 = [0 :_R (0 :_R I_1)] \cap [0 :_R (0 :_R I_2)] \\ &= 0 :_R (0 :_R I_1 + 0 :_R I_2). \end{aligned}$$

So

$$\begin{aligned} 0 :_R (I_1 \cap I_2) &= 0 :_R [0 :_R (0 :_R (I_1 \cap I_2))] = 0 :_R [0 :_R (0 :_R I_1 + 0 :_R I_2)] \\ &= 0 :_R I_1 + 0 :_R I_2. \end{aligned}$$

By [5, Theorem 1] R is self FP-injective.

(\Leftarrow) Let M be a finitely presented R -module and let $0 \neq x \in M$. We claim that there is a nonzero homomorphism $h : Rx \rightarrow R$ with $h(x) \neq 0$. Otherwise, suppose $(Rx)^* = \text{Hom}_R(Rx, R) = 0$. Since Rx is finitely presented, there is an exact sequence

$F_1 \rightarrow F_0 \rightarrow Rx \rightarrow 0$ with F_0 and F_1 finitely generated free modules. Then $0 \rightarrow F_0^* \rightarrow F_1^* \rightarrow A \rightarrow 0$ is exact where $A = \text{Coker}(F_0^* \rightarrow F_1^*)$. Consider the following commutative diagram with exact rows.

$$\begin{array}{ccccccc} F_1 & \longrightarrow & F_0 & \longrightarrow & Rx & \longrightarrow & 0 \\ \downarrow \sigma_{F_1} & & \downarrow \sigma_{F_0} & & \downarrow \varphi & & \\ F_1^{**} & \longrightarrow & F_0^{**} & \longrightarrow & \text{Ext}_R^1(A, R) & \longrightarrow & 0 \end{array}$$

where $\sigma_{F_0}, \sigma_{F_1}$ are the canonical evaluation homomorphisms, φ is an induced homomorphism. It is known that $\sigma_{F_0}, \sigma_{F_1}$ are isomorphisms, so φ is also an isomorphism and hence $\text{Ext}_R^1(A, R) \cong Rx \neq 0$, which contradicts that R is self FP-injective since A is finitely presented.

Since Rx and M are finitely presented, a nonzero homomorphism $h : Rx \rightarrow R$ can be extended to a homomorphism $\bar{h} : M \rightarrow R$ with $\bar{h}(x) = h(x) \neq 0$. Thus $\text{Rej}_M(R) = 0$, and R cogenerates M by [1, Corollary 8.13]. The proof is finished. \square

We now in a position to give the main result.

Theorem 1 *The following statements are equivalent.*

- (1) R is a QF ring;
- (2) R is an artin ring and R^e is flat;
- (2') R is a noether ring and R^e is flat;
- (3) M^e is a projective module for any free (projective, flat) module M ;
- (3') M^e is a submodule of a projective module for any free (projective, flat) module M ;
- (4) R is an E -artin ring and R is self FP-injective;
- (5) R is an artin ring and R is an f -cogenerator in the category of R -modules;
- (5') R is a noether ring and R is an f -cogenerator in the category of R -modules;
- (6) R is an artin ring and some injective cogenerator is flat;
- (6') R is a noether ring and some injective cogenerator is flat;
- (7) R is an artin ring and $E(R/m)$ is flat for each $m \in \text{Max}(R)$, where $\text{Max}(R)$ is the maximal spectrum of R ;
- (7') R is a noether ring and $E(R/m)$ is flat for each $m \in \text{Max}(R)$.

Proof (1) \Leftrightarrow (2) \Leftrightarrow (2') \Leftrightarrow (3) \Leftrightarrow (3') See Proposition 1.

(1) \Leftrightarrow (5) \Leftrightarrow (5') follow easily from Lemma 4.

(1) \Rightarrow (4), (1) \Rightarrow (6) \Rightarrow (6') \Rightarrow (7)' and (1) \Rightarrow (7) \Rightarrow (7)' are trivial.

(4) \Rightarrow (1) Suppose that Q is an injective module. We know that Q is a submodule of E^I for some set I . Since R is self FP-injective, $R^{(I)}$ is FP-injective. By Lemma 2 E^I is a submodule of a projective module P because $E^I \cong [R^{(I)}]^e$. It follows that Q is a submodule of P . Hence Q is projective and then R is a QF ring.

(7)' \Rightarrow (1) Suppose that R is a noether ring and $E(R/m)$ is flat for each $m \in \text{Max}(R)$. Let $E_1 = \bigoplus_{m \in \text{Max}(R)} E(R/m)$. Then E_1 is flat. It follows from [8, Theorem 9.51] that $\text{Hom}_R(\text{Ext}_R^1(R/I, R), E_1) \cong \text{Tor}_1^R(\text{Hom}_R(R, E_1), R/I)$ for any ideal I of R . Since

$$\text{Tor}_1^R(\text{Hom}_R(R, E_1), R/I) \cong \text{Tor}_1^R(E_1, R/I) = 0,$$

$$\text{Hom}_R(\text{Ext}_R^1(R/I, R), E_1) = 0.$$

It is known [1, Corollary 18.16] that E_1 is an injective cogenerator in the category of R -modules, so $\text{Ext}_R^1(R/I, R) = 0$ and hence R is selfinjective and R is a QF ring. \square

Definition 3 R is called an E -coherent ring if Q^e is a submodule of a flat module for any injective module Q .

Remark 3 By [4, Theorem 1], a coherent ring is an E -coherent ring. If R is a semihereditary ring, then R is coherent if and only if R is E -coherent.

Remark 4 We can get similar conclusions about E -coherent rings to that about E -artin rings in Lemmas 1 and 2, which we omit.

Theorem 2 The following statements are equivalent.

- (1) R is an IF ring;
- (2) R is a coherent ring and R^e is flat;
- (3) M^e is a flat module for any free (projective, flat) module M ;
- (3') M^e is a submodule of a flat module for any free (projective, flat) module M ;
- (4) R is an E -coherent ring and R is self FP-injective;
- (5) R is a coherent ring and R is an f -cogenerator in the category of R -modules;
- (6) R is a coherent ring and some injective cogenerator is flat;
- (7) R is a coherent ring and $E(R/m)$ is flat for each $m \in \text{Max}(R)$.

Proof The proof is similar to that of Theorem 1, and is omitted here. \square

Theorem 3 Consider the following conditions.

- (1) R is a semihereditary ring;
- (2) M^e is an FP-injective module for any finitely presented module M ;
- (3) M^{ee} is a flat module for any finitely presented module M ;
- (4) M^{eee} is an FP-injective module for any finitely presented module M .

In general (1) \Leftarrow (2) \Leftrightarrow (3) \Leftrightarrow (4). If R is self FP-injective, then the above conditions are equivalent.

Proof (2) \Rightarrow (1) Suppose K is a finitely generated submodule of a projective module. Then K is a submodule of some finitely generated free module R^n . So we have an exact sequence $0 \rightarrow K \rightarrow R^n \rightarrow M \rightarrow 0$ with M finitely presented, and hence $0 \rightarrow M^e \rightarrow (R^n)^e \rightarrow K^e \rightarrow 0$ is exact. Since M^e is FP-injective by (2), the latter exact sequence is pure. So it splits by [4, Lemma 1]. Because $(R^n)^e \cong E^n$ is injective, M^e is also injective. Then M is flat by [6, Theorem 1.4]. It follows from [8, Theorem 3.57] that $0 \rightarrow K \rightarrow R^n \rightarrow M \rightarrow 0$ splits. Thus K is projective and R is a semihereditary ring.

(2) \Rightarrow (3) If (2) holds, then R is a semihereditary ring. So R is a coherent ring. If M is a finitely presented module, then M^e is FP-injective by (2). It follows from [4, Theorem 1] that M^{ee} is flat.

(3) \Rightarrow (4) It follows from [6, Theorem 1.4].

(4) \Rightarrow (2) Suppose M is a finitely presented module. Then M^{eee} is FP-injective by (4). Since M^e is a direct summand of M^{eee} by [9, Exercise 23, p. 46], M^e is FP-injective.

Now let R be self FP-injective. We will show that (1) implies (2).

Let R be a semihereditary ring. Since R is self FP-injective, R is an IF ring. Suppose that M is a finitely presented module. Then there is an exact sequence $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ with K a finitely generated module and F a finitely generated projective module. K^e is flat by Theorem 2, so the exact sequence $0 \rightarrow M^e \rightarrow F^e \rightarrow K^e \rightarrow 0$ is pure. It follows that M^e is a pure submodule of the injective module F^e . So M^e is FP-injective. This completes the proof of this theorem. \square

References:

- [1] ANDERSON F W, FULLER K R. *Rings and Categories of Modules* [M]. 2nd ed, Graduate Texts in Mathematics, Vol. 13, Springer-Verlag, 1992.
- [2] CARTAN H, EILENBERG S. *Homological Algebra* [M]. Princeton University Press, 1956.
- [3] GLAZ S. *Commutative Coherent Rings* [M]. Lecture Notes in Mathematics, Vol. 1371, Springer-Verlag, 1989.
- [4] HUANG Z Y. *On the flatness and injectivity of dual modules* [J]. Southeast Asian Bull. of Math., 1997, 21: 257-262.
- [5] IDEDA M, NAKAYAMA T. *On some characteristic properties* [J]. Proc. Amer. Math. Soc., 1954, 5: 15-19.
- [6] ISHIKAWA T. *On injective and flat modules* [J]. Jour. Math. Soc. Japan, 1965, 17: 291-296.
- [7] JAIN S. *Flat and FP-injectivity* [J], Proc. Amer. Math. Soc., 1973, 41: 437-442.
- [8] ROTMAN J J. *An Introduction to Homological Algebra* [M]. Academic Press, New York, 1979.
- [9] STENTRÖM B. *Rings of Quotients* [M], Spinger-Verlag, Springer, 1975.

关于对偶模的平坦性和内射性 (II)

黄兆泳¹, 唐金玉²

(1. 南京大学数学系, 江苏 南京 210093; 2. 广西大学数学与信息科学系, 广西 南宁 530004)

摘要: 对交换环 R 和 R -模范畴上的一个内射余生成元 E , 我们用相对于 E 的对偶模的性质刻画了 QF 环, IF 环和半遗传环.